arXiv:1210.5661v1 [cond-mat.stat-mech] 20 Oct 2012

Hardwiring a Maxwell demon

Philipp Strasberg[1], Gernot Schaller[1], Tobias Brandes[1], and Massimiliano Esposito[2]
[1] *Institut für Theoretische Physik, Technische Universität Berlin, Hardenbergstr. 36, D-10623 Berlin, Germany*
[2] *Complex Systems and Statistical Mechanics, University of Luxembourg, L-1511 Luxembourg, Luxembourg*

We present a physical implementation of a Maxwell demon which consists of a conventional single electron transistor (SET) capacitively coupled to another quantum dot detecting its state. Altogether, the system is described by stochastic thermodynamics. We identify the regime where the energetics of the SET is not affected by the detection, but where its coarse-grained entropy production is shown to contain a new contribution compared to the isolated SET. This additional contribution can be identified as the information flow generated by the "Maxwell demon" feedback in an idealized limit.

PACS numbers: 05.70.Ln, 05.40.-a 05.60.Gg,

It has been more than a century that the thermodynamic implications of various types of "intelligent interventions" (e.g. feedbacks) on the microscopic degrees of freedom of a system have intrigued scientists [1]. A Maxwell demon for example can be thought of as a hidden idealized mechanism which is able to modify the second law of thermodynamics (entropy balance) but without modifying the first law (energy balance). Such a demon would thus be able to heat a hot reservoir while cooling down a cold reservoir without using any additional energy, which clearly breaks the traditional formulation of the second law of thermodynamics.

Nowadays, our ability to manipulate small devices has drastically increased, and what used to be unrealistic thought experiments have become real experiments [2–4]. In parallel to that, significant progress in understanding the nonequilibrium thermodynamics of small systems has been achieved [5–7]. This is particularly true for systems described by Markovian stochastic dynamics where a consistent theoretical framework, called stochastic thermodynamics, has emerged and has been shown very useful to study fluctuations and efficiencies of systems driven far from equilibrium [8–10]. Quite naturally, recent studies have started considering the thermodynamic description of systems subjected to different types of feedbacks [11–24].

To understand the thermodynamic behaviour of feedback controlled systems it is important to include the *information* generated or used by the feedback [22, 24]. Nevertheless, every feedback scheme has to be implemented physically and the natural question which arises – and which has not been answered yet – is under which circumstances this physical implementation appears as pure information entering the thermodynamic description. This is the object of the present letter.



FIG. 1: (Color online) The shaded region constitutes the demon, the contained dot d couples capacitively to the lower dot s via Coulomb repulsion U. The latter is placed in a conventional SET-setup, which can be accessed experimentally, whereas the demon remains hidden. The trajectory (with initial state $\rho = 1$ and $\sigma = E$) marked by the steps $1-4$, where an electron is transfered against the bias through the SET, becomes likely in the Maxwell demon limit.

Our model consists of two single level quantum dots interacting capacitively via a Coulomb repulsion U and additionally coupled to thermal reservoirs as depicted in Fig. 1. No electrons can be transfered between the dots. The corresponding system Hamiltonian is given by $H = \epsilon_d c_d^\dagger c_d + \epsilon_s c_s^\dagger c_s + U c_d^\dagger c_d c_s^\dagger c_s$ with fermionic operators annihililating electrons on the system dot (c_s) and the detecting dot (c_d). The dots are weakly coupled to ideal reservoirs $\nu \in \{D, L, R\}$ at temperature $T_\nu = 1/\beta_\nu$ ($k_B \equiv 1$) and chemical potential μ_ν. As shown in Fig. 1, dot s is coupled to the two reservoirs L and R and constitutes a usual SET, while dot d is coupled to reservoir D. One can show that dot d with reservoir D can be tuned to "detect" the state of the SET [25] and will eventually constitute the Maxwell demon. We denote the four eigenvectors of the two coupled dots by $|\rho\sigma\rangle = |\rho\rangle_d \otimes |\sigma\rangle_s$ where $\rho \in \{0, 1\}$ and $\sigma \in \{E, F\}$ denote the states of the dot d and s, respectively, which are either empty (0 or E) or filled (1 or F). For weak dot-reservoir interaction, the time evolution of the coupled dots density matrix ρ can be shown to be governed by a Markovian master equation $\frac{d}{dt}\rho = \mathcal{W}\rho$, which simply yields a rate equation for the probabilities $p_{\rho\sigma}$ to be in the eigenstate $|\rho\sigma\rangle$. In the ordered basis $(p_{0E}, p_{1E}, p_{0F}, p_{1F})$ the rate matrix reads

$$\mathcal{W} = \begin{pmatrix} \mathcal{W}_{EE} & \mathcal{W}_{EF} \\ \mathcal{W}_{FE} & \mathcal{W}_{FF} \end{pmatrix} \equiv \sum_\nu \mathcal{W}^{(\nu)} . \tag{1}$$

The two blocks

$$\mathcal{W}_{EE} = \begin{pmatrix} -\gamma_D - \gamma_L - \gamma_R & \overline{\gamma}_D \\ \gamma_D & -\overline{\gamma}_D - \gamma_L^U - \gamma_R^U \end{pmatrix} , \tag{2}$$

$$\mathcal{W}_{FF} = \begin{pmatrix} -\gamma_D^U - \overline{\gamma}_L - \overline{\gamma}_R & \overline{\gamma}_D^U \\ \gamma_D^U & -\overline{\gamma}_D^U - \overline{\gamma}_L^U - \overline{\gamma}_R^U \end{pmatrix} \tag{3}$$

denote the dynamics of the dot d when dot s is empty or filled respectively, while the blocks

$$\mathcal{W}_{EF} = \mathrm{Diag}(\overline{\gamma}_L + \overline{\gamma}_R, \overline{\gamma}_L^U + \overline{\gamma}_R^U) , \tag{4}$$

$$\mathcal{W}_{FE} = \mathrm{Diag}(\gamma_L + \gamma_R, \gamma_L^U + \gamma_R^U) \tag{5}$$

denote the transitions between an empty and filled dot s which can occur when dot d is either empty or filled. The rates are given by $\gamma_\nu \equiv \Gamma_\nu f_\nu$, $\gamma_\nu^U \equiv \Gamma_\nu^U f_\nu^U$, $\overline{\gamma}_\nu \equiv \Gamma_\nu(1 - f_\nu)$ and $\overline{\gamma}_\nu^U \equiv \Gamma_\nu^U(1 - f_\nu^U)$ with electronic tunneling rates $\Gamma_\nu > 0$ and $\Gamma_\nu^U > 0$ and Fermi functions $0 < f_\nu, f_\nu^U < 1$ with $\nu \in \{D, L, R\}$. Note that $\Gamma_\nu \neq \Gamma_\nu^U$ requires to go beyond the common wide-band approximation. The Fermi functions are evaluated at the respective transition energies of the SET dot $f_{L/R} \equiv 1/[\exp(\beta_{L/R}(\epsilon_s - \mu_{L/R})) + 1]$, $f_{L/R}^U \equiv 1/[\exp(\beta_{L/R}(\epsilon_s + U - \mu_{L/R})) + 1]$ and the detector dot $f_D \equiv 1/[\exp(\beta_D(\epsilon_d - \mu_D)) + 1]$, $f_D^U \equiv 1/[\exp(\beta_D(\epsilon_d + U - \mu_D)) + 1]$, respectively.

This system is consistently described by stochastic thermodynamics. At steady state the entropy production of the full system reads

$$\dot{S}_i = \sum_\nu \sum_{\rho,\rho',\sigma,\sigma'} W^{(\nu)}_{\rho\sigma\rho'\sigma'} p_{\rho'\sigma'} \ln \frac{W^{(\nu)}_{\rho\sigma\rho'\sigma'}}{W^{(\nu)}_{\rho'\sigma'\rho\sigma}} \geq 0 \tag{6}$$

and can be rewritten after some algebra as the sum of force-flux terms associated to matter and energy transfers, such that e.g. for $\beta_L = \beta_R = \beta$ we obtain

$$\dot{S}_i = \beta(\mu_L - \mu_R) I_S + (\beta_D - \beta) I_E \geq 0, \tag{7}$$

where $I_S = \gamma_L p_{0E} - \overline{\gamma}_L p_{0F} + \gamma_L^U p_{1E} - \overline{\gamma}_L^U p_{1F}$ is the stationary electronic particle current flowing from reservoir L to R through the SET and $I_E = U(\gamma_D p_{0E} - \overline{\gamma}_D p_{1E})$ is the energy current entering the reservoir D due to the interaction between dot d and the SET. The matter current associated to the detector bath D vanishes since there is no particle exchange between dot d and dot s.

Using usual techniques [6, 26], it is possible to show that the following fluctuation theorem for the entropy production is satisfied:

$$\lim_{t\to\infty} \frac{p_{+n_S,+n_D}(t)}{p_{-n_S,-n_D}(t)} = e^{\beta(\mu_L - \mu_R) n_S + (\beta_D - \beta) U n_D} , \tag{8}$$

where $p_{n_S,n_D}(t)$ denotes the probability of having n_S electrons traversing the system from left to right together with net n_D electrons entering dot d at energy ϵ_d and leaving it at energy $\epsilon_d + U$, altogether leading to a net energy transfer of $U n_D$ into reservoir D after time t. So far, our system is thus only a conventional thermoelectric device in which the thermal gradient may be used to generate an electronic current through the SET against the bias. Related models have been considered in [25, 27–30].

We now assume that our experimental setup allows us to detect electron transfers in the SET (e.g. counting statistics experiments) but does not provide any information about the existence of the demon (i.e. dot d and reservoir D). The observed SET states $\sigma \in \{E, F\}$ thus constitute two coarse-grained “mesostates” with probabilities $p_\sigma = p_{0\sigma} + p_{1\sigma}$, see Fig. 2. Moreover, we denote by $\mathbb{P}_{\rho|\sigma} = p_{\rho\sigma}/p_\sigma$ the conditional stationary probability to find dot d in state ρ if the state of the SET is σ. Each of these probabilities can be explicitly calculated from the analytical steady-state solution of Eq. (1). It is straightforward to see that the exact coarse-grained dynamics of the SET may formally be written as



FIG. 2: (Color online) Visualization of the dynamics in the full (black arrows) and coarse-grained (red arrows) state space. The four states are denoted by black squares with their size being proportional to the occupation probability near the sought-after limit. The two mesostates of the SET (E or F) are composed by the shaded regions. They each contain two possible states of the detector dot d which are connected by fast transitions. The red labels denote transition rates between the two SET mesostates in the error free limit ($f_D \to 1$, $f_D^U \to 0$ and $\Gamma_D/\Gamma_{L/R} \to \infty$).

$$\dot{p}_\sigma = \sum_{\sigma'} V_{\sigma\sigma'} p_{\sigma'} \tag{9}$$

with ”rates” $V_{\sigma\sigma'} = \sum_\nu V^{(\nu)}_{\sigma\sigma'} = \sum_\nu \sum_{\rho\rho'} W^{(\nu)}_{\rho\sigma,\rho'\sigma'} \mathbb{P}_{\rho'|\sigma'}$.

Fast demon: Since we want dot d and reservoir D to ultimately constitute a Maxwell demon, we are now going to assume that the dynamics of the demon is much faster than the SET dynamics $\Gamma_D = \Gamma_D^U \gg \max\{\Gamma_L^{(U)}, \Gamma_R^{(U)}\}$. As expected, in the extreme case $\Gamma_D/\max\{\Gamma_{L/R}^{(U)}\} \to \infty$, the conditional probabilities equilibrate instantaneously with respect to the reservoir D

$$\begin{aligned} \mathbb{P}_{0|E} &\to \frac{\overline{\gamma}_D}{\gamma_D + \overline{\gamma}_D} = 1 - f_D\,, \quad \mathbb{P}_{1|E} \to 1 - \mathbb{P}_{0|E}\,, \\ \mathbb{P}_{0|F} &\to \frac{\overline{\gamma}_D^U}{\gamma_D^U + \overline{\gamma}_D^U} = 1 - f_D^U\,, \quad \mathbb{P}_{1|F} \to 1 - \mathbb{P}_{0|F}\,, \end{aligned} \tag{10}$$

such that Eq. (9) becomes an ordinary rate equation. For finite demon temperatures, there will thus always be some finite detection "error", which we quantify by $\epsilon_E \equiv \mathbb{P}_{0|E}/\mathbb{P}_{1|E}$ and $\epsilon_F \equiv \mathbb{P}_{1|F}/\mathbb{P}_{0|F}$. For instance, when $\epsilon_d = \mu_D - U/2$, these errors will be given by $\epsilon_E = \epsilon_F = \exp(-\beta_D U/2)$.

The entropy production corresponding to this coarse-grained SET dynamics is

$$\dot{S}_i^* = \sum_{\nu,\sigma,\sigma'} V_{\sigma\sigma'}^{(\nu)} p_{\sigma'} \ln \frac{V_{\sigma\sigma'}^{(\nu)}}{V_{\sigma'\sigma}^{(\nu)}} = A_S^* I_S \geq 0 \tag{11}$$

with an effective affinity

$$A_S^* = \ln\left[\frac{\left(\overline{\gamma}_D \gamma_L + \gamma_D \gamma_L^U\right)\left(\overline{\gamma}_D^U \overline{\gamma}_R + \gamma_D^U \overline{\gamma}_R^U\right)}{\left(\overline{\gamma}_D^U \overline{\gamma}_L + \gamma_D^U \overline{\gamma}_L^U\right)\left(\overline{\gamma}_D \gamma_R + \gamma_D \gamma_R^U\right)}\right]. \tag{12}$$

The coarse-grained entropy production can be shown to always under-evaluate the full entropy production: $\Delta\dot{S}_i \equiv \dot{S}_i - \dot{S}_i^* \geq 0$ [31]. Furthermore, it implies an effective fluctuation theorem for the particle counting statistics of the SET (e.g. see [30])

$$\lim_{t\to\infty} \frac{p_{+n_S}(t)}{p_{-n_S}(t)} = e^{A_S^* n_S}\,, \tag{13}$$

where $p_{n_S}(t)$ is the probability of having n_S electrons transferred from left to right after time t. This demonstrates that the coarse-grained entropy production (11) is a meaningful and measurable quantity in the fast-demon limit. Subtracting the entropy production of an isolated SET from the coarse-grained entropy production yields an additional term

$$I_F \equiv \dot{S}_i^* - \dot{S}_i^{\rm SET} = \dot{S}_i^* - \ln\left[\frac{f_L(1-f_R)}{(1-f_L)f_R}\right] I_S\,, \tag{14}$$

which in ignorance of the physical nature of the demon must be interpreted as an information current between demon and system.

Fast and Precise demon: Since the demon also needs to be able to reliably discriminate between the two states of the SET, we put $\epsilon_d = \mu_D - U/2$ and further assume that $\beta_D U \gg 1$, which implies $\overline{\gamma}_D, \gamma_D^U \to 0$ and $\gamma_D, \overline{\gamma}_D^U \to \Gamma_D$. Then, the two dots become perfectly correlated and the steady state reads $p_{1E} = (\overline{\gamma}_L + \overline{\gamma}_R)/(\overline{\gamma}_L + \overline{\gamma}_R + \gamma_L^U + \gamma_R^U)$, $p_{0F} = 1 - p_{1E}$, $p_{0E} = p_{1F} = 0$. The effective affinity (12) can in this limit be written as

$$A_S^* = \ln\left[\frac{f_L(1-f_R)}{(1-f_L)f_R} \frac{f_L^U f_R}{f_L f_R^U} \frac{\Gamma_L^U \Gamma_R}{\Gamma_L \Gamma_R^U}\right], \tag{15}$$

which implies for the information current

$$I_F = \ln\left[\frac{f_L^U f_R}{f_L f_R^U} \frac{\Gamma_L^U \Gamma_R}{\Gamma_L \Gamma_R^U}\right] I_S\,. \tag{16}$$

This contribution results from the interaction with dot d which we have traced out. We note however, that this demon is effectively extracting energy from the SET at a rate

$$I_E = U \frac{(\overline{\gamma}_L + \overline{\gamma}_R)(\gamma_L^U + \gamma_R^U)}{\overline{\gamma}_L + \overline{\gamma}_R + \gamma_L^U + \gamma_R^U}. \tag{17}$$

and is thus creating an imbalance between the energy currents at the left and right interface of the SET. It is therefore not yet a true Maxwell demon. In fact, the right hand side equals $U/2$ times the *activity current* in the SET which measures the total number of electron jumps in and out of the SET. This is due to the fact that each change in the mesostates ($E \leftrightarrow F$) instantaneously induces a jump in dot d ($1 \leftrightarrow 0$) and in the error-free limit no other contribution arises. Finally, we note that due to the finite energy current I_E, the entropy flow between dot d and its reservoir, $\dot{S}_e^{(D)} = -\beta_D I_E$ must diverge as $\beta_D \to \infty$, which also holds for the total entropy production $\dot{S}_i = -\sum_\nu \dot{S}_e^{(\nu)}$.

Maxwell demon: In order to obtain a true Maxwell demon, we further need to assume that the temperatures of the left and right reservoirs are sufficiently large compared to the capacitive interaction U such that $\beta U \to 0$. In this limit, the energetics of the SET is not affected anymore by the demon since the Fermi functions evaluated at the different energies become equal:

$$\lim_{\beta U\to 0} f_\nu^U = f_\nu \quad , \quad \nu \in \{L, R\}. \tag{18}$$

A naive experimenter assuming tight coupling between energy and particle SET currents would therefore conclude conservation of energy flowing into the system from left and right terminals. In contrast, a suspicious experimenter with the ability to monitor both SET energy currents would still observe that energy currents into left and right reservoirs are not conserved $I_E^{(L)} + I_E^{(R)} = -I_E$ to order U, cf. Eq. (17). In comparison to the SET energy current at e.g. the right lead, this is a correction of order U/ϵ_s, which can be made arbitrarily small nearly everywhere.

The bare rates Γ_ν could and – to obtain any nontrivial effect – should however depend on the energy levels of the dots. As a result, the information current (16) becomes

$$I_F = (\delta_L - \delta_R) I_S, \tag{19}$$

where we introduced the feedback parameters $\delta_\nu = \ln\left[\Gamma_\nu^U/\Gamma_\nu\right]$ with $\nu \in \{L, R\}$. Remarkably, the coarse-grained rate matrix (9) describing the effective SET dynamics now satisfies the modified local detailed balance condition

$$\ln \frac{V_{FE}^{(\nu)}}{V_{EF}^{(\nu)}} = -\beta_\nu(\epsilon_s - \mu_\nu) + \delta_\nu \quad , \quad \nu \in \{L, R\}. \tag{20}$$

This result is in perfect agreement with the modified local detailed balance condition introduced in [24] to describe Maxwell demon feedbacks within the framework of stochastic thermodynamics. Alternatively, such a modification of local detailed balance may be generated using a fast feedback control loop [32]. Now, when $\delta_L \ll 0$ and/or $\delta_R \gg 0$, a trajectory as shown in Fig. 1 becomes highly probable. Naturally, even without the strict mathematical limits that we discussed, we demonstrate in Fig. 3 that the coarse-grained entropy production $\dot{S}_i^*$ approaches the ideal Maxwell demon from [24] in Fig. 3. Furthermore, the inset demonstrates that for $\epsilon_s \gg U$, the modification of the first law for an isolated SET $I_E^{(L)} + I_E^{(R)} = -I_E \approx 0$ is negligible.



FIG. 3: (Color online) The coarse-grained entropy production $\dot{S}_i^*$ as a function of the SET bias βV in the fast demon limit (solid brown curves) differs from the unperturbed SET case (dashed magenta) in non-equilibrium and moves towards the ideal Maxwell demon limit (dashed green) as the temperature of the detector bath (and correspondingly the error of the demon) is lowered ($\beta_D = 1\beta, 200\beta, 800\beta$ with increasing thickness, solid arrow). The total entropy production always exceeds the coarse-grained one for similar parameters (e.g. dash-dotted blue vs. bold-solid brown for $\beta_D = 800$). Dotted blue curves demonstrate the rapid convergence of the full entropy production to the fast demon limit when $\Gamma_D \gg \Gamma$ (dotted arrow). The SET (solid) and demon (dotted) energy current in the inset (for $\Gamma_D = 100\Gamma$ and $\beta_D = 800\beta$) demonstrate that the relative modification of the SET's first law is of order U/ϵ_s, which can be made arbitrarily small in comparison to the SET energy currents nearly everywhere. Other parameters were chosen as $\Gamma_L = \Gamma_R^U = e^{+\delta}\Gamma$, $\Gamma_L^U = \Gamma_R = e^{-\delta}\Gamma$ with $\delta = \ln 2$, $\beta_L = \beta_R = \beta$, $\mu_L = \epsilon_s + V/2$, $\mu_R = \epsilon_s - V/2$, $\beta\epsilon_s = 1$, $\beta U = 0.01$.

Finally, we note that using the coarse-grained entropy production (11), a negative information current can be used to define various efficiencies (e.g. using information to transport electrons against the bias or to cool a cold reservoir). In contrast, the efficiencies of the thermoelectric device as a whole may be drastically different, as follows from the divergence of the total entropy production in the fast and precise demon limit.

Conclusion To the best of our knowledge, this letter establishes for the first time the precise connection between the complete thermodynamic description of a Maxwell demon model and the system it is acting on. In particular, we have identified the effective level of description of the system where the demon manifests itself solely through an information flow modifying the second law. Furthermore, by showing that the effective entropy production is only a piece of the total entropy production of the joined system, we provide a rigorous support for the generic claim that a system subjected to a "Maxwell demon" is an idealization which neglects the dissipation associated to the implementation of the demon mechanism.

Acknowledgments Financial support by the DFG (SCHA 1646/2-1, SFB 910, and GRK 1558) and the National Research Fund, Luxembourg (project FNR/A11/02) is gratefully acknowledged.

---

[1] H. Leff and A. F. Rex, *Maxwell's Demon 2: Entropy, Classical and Quantum Information, Computing* (CRC Press, 2002).

[2] V. Serreli, C.-F. Lee, E. R. Kay, and D. A. Leigh, Nature **445**, 523 (2007).

[3] S. Toyabe, T. Sagawa, M. Ueda, E. Muneyuki, and M. Sano, Nature Physics **6**, 988 (2010).

[4] A. Brut, A. Arakelyan, A. Petrosyan, S. Ciliberto, R. Dillenschneider, and E. Lutz, Nature **183**, 487 (2012).

[5] C. Jarzynski, Annual Review of Condensed Matter Physics **2**, 329 (2011).

[6] M. Esposito, U. Harbola, and S. Mukamel, Rev. Mod. Phys. **81**, 1665 (2009).

[7] M. Campisi, P. Hänggi, and P. Talkner, Rev. Mod. Phys. **83**, 771 (2011).

[8] U. Seifert, arXiv:1205.4176 (2012).

[9] M. Esposito and C. Van den Broeck, Phys. Rev. E **82**, 011143 (2010).

[10] C. Van den Broeck and M. Esposito, Phys. Rev. E **82**, 011144 (2010).

[11] R. Kawai, J. M. R. Parrondo, and C. Van den Broeck, Phys. Rev. Lett. **98**, 080602 (2007).

[12] T. Sagawa and M. Ueda, Phys. Rev. Lett. **100**, 080403 (2008).

[13] T. Sagawa and M. Ueda, Phys. Rev. Lett. **102**, 250602 (2009).

[14] T. Sagawa and M. Ueda, Phys. Rev. Lett. **104**, 090602

(2010).
[15] Y. Zhou and D. Segal, Phys. Rev. E **82**, 011120 (2010).
[16] J. M. Horowitz and S. Vaikuntanathan, Phys. Rev. E **82**, 061120 (2010).
[17] J. Horowitz and J. M. P. Parrondo, EPL **95**, 10005 (2011).
[18] J. Horowitz and J. M. P. Parrondo, New J. Phys. **13**, 123019 (2011).
[19] M. Esposito and C. Van den Broeck, EPL **95**, 40004 (2011).
[20] D. Abreu and U. Seifert, EPL **94**, 10001 (2011).
[21] M. Bauer, D. Abreu, and U. Seifert, J. Phys. A **45**, 162001 (2012).
[22] T. Sagawa and M. Ueda, arXiv:1206.2479 (2012).
[23] D. Mandal and C. Jarzynski, PNAS **109**, 11641 (2012).
[24] M. Esposito and G. Schaller, EPL **99**, 30003 (2012).
[25] G. Schaller, G. Kießlich, and T. Brandes, Phys. Rev. B **82**, 041303 (2010).
[26] D. Andrieux and P. Gaspard, J. Stat. Phys. **127**, 107 (2007).
[27] B. Rutten, M. Esposito, and B. Cleuren, Phys. Rev. B **80**, 235122 (2009).
[28] R. Sanchez, R. Lopez, D. Sanchez, and M. Buttiker, Phys. Rev. Lett. **104**, 076801 (2010).
[29] R. Sánchez and M. Büttiker, Phys. Rev. B **83**, 085428 (2011).
[30] G. Bulnes Cuetara, M. Esposito, and P. Gaspard, Phys. Rev. B **84**, 165114 (2011).
[31] M. Esposito, Phys. Rev. E **85**, 041125 (2012).
[32] G. Schaller, C. Emary, G. Kießlich, and T. Brandes, Phys. Rev. B. **84**, 085418 (2011).